

# Health Measurement in the Third Era of Health

When writing about “the second epidemiologic revolution,” Terris discussed 2 eras in health. The first era—the communicable disease era—began during ancient times and continues today; the second era—the chronic disease era—began during the 20th century, particularly among the industrialized nations.

Although neither revolution against these types of diseases is complete, we have made such considerable progress that substantial and growing segments of the population no longer regard disease as the only, or even the primary, health problem. Increasingly, the goal is a long and fruitful life, not simply the absence of disease. That potential and the effort to achieve it compose the third era of health, and a proposed new measure of health is outlined in this article. (*Am J Public Health*. 2006;96:17–19. doi:10.2105/AJPH.2004.055970)

Lester Breslow, MD, MPH

**UNTIL RECENT TIMES,** health measurement was appropriately focused on disease and injury and their consequences, particularly disability and death, and the activities intended to alleviate disease and injury, particularly medical and related services. However, strictly speaking, those services are not a part of health measurement. During the evolution of mankind, biological susceptibility to disease and injury has been carried forward with the environmental conditions that induce health impairments. Health measurement has increasingly been extended to include these important associations with health.

## TWO HEALTH REVOLUTIONS

Terris wrote about 2 eras, or revolutions, in health.<sup>1</sup> The first health revolution was against communicable diseases, and that revolution continues today. The second revolution, which began in the mid-20th century and also continues today, was against chronic diseases. Whereas most health professionals who were concerned with the second revolution focused on institutional care for the affected, epidemiologists began to examine the causative factors of these newly prominent diseases, which they regarded as epidemics that have lasted for decades rather than weeks or months. Their scientific investigations showed how tobacco, excessive fat consumption, lack of exercise, and other

factors characteristic of life in industrialized countries during the 20th century caused the chronic disease epidemic.<sup>2,3</sup>

The so-called *epidemiological transition*—from communicable diseases to chronic diseases as the primary public health problem—first occurred among the well-to-do in developed countries, because they were the first to be exposed to the causative conditions. The epidemic of chronic disease then moved down the social hierarchy as the less affluent were exposed to the causative conditions. During the 20th century, coronary heart disease, lung cancer, and other chronic diseases were mass phenomena. However, during the past 50 years, considerable progress has been made against the chronic diseases with the second health revolution, which is well underway in the industrialized world but is only beginning in developing countries. For example, heart disease mortality in the United States has been declining since 1950, and cancer mortality reached a peak in 1990 before it began to decline (Table 1).

## THE THIRD ERA OF HEALTH

With the first and second health revolutions so far advanced, what is next? Actually, we have already entered the third era of health,<sup>6</sup> a time when people are living into their 70s and 80s and are increasingly free of disease burdens.

Ninety percent of all Americans, and almost 70% of those older than 75 years, believe their health is excellent or good, not just fair or poor (Table 2). People now seek to develop and maintain their health, not merely combat disease, which reflects the progress against both communicable and chronic diseases. All life activities require a certain anatomical, physiological, sensory, mental, or other health competence—a personal resource. The World Health Organization's (WHO) Ottawa Charter thus defined health as “a resource for everyday life.”<sup>7</sup>

In 1948, the WHO said health was a positive notion and that it was “physical, mental, and social well-being, not merely the absence of disease and infirmity.”<sup>8</sup> Efforts have been made

**TABLE 1—Gross Death Rates per 100 000 From Major Chronic Diseases: United States, 1900–2000**

	1900	1950	2000
Heart Disease	137	356	253
Cancer	64	140	197
Cerebrovascular disease	107	104	60

Sources: National Office of Vital Statistics<sup>4</sup> and National Center for Health Statistics.<sup>5</sup>

**TABLE 2—Self-Assessed Health Status as Excellent or Good: United States, 1991–2001**

	1991	1995	2000	2001
Total, %	89.6	89.4	91.0	90.8
Age, y				
<18, %	97.4	97.4	98.3	98.2
18–44, %	93.9	93.4	94.9	94.6
45–54, %	86.6	86.6	88.1	88.3
55–64, %	79.3	78.6	82.1	80.8
≥65, %	71.0	71.7	73.0	73.4
≥75, %	66.4	67.8	67.8	69.2

Source: National Center for Health Statistics.<sup>5</sup>

to measure health with that WHO concept.<sup>9–13</sup> One such effort noted that health status can be delineated as a spectrum (excellent to very poor) on which every person can be placed at any one time.<sup>13</sup> In that early formulation, however, health was a state of well-being. Going beyond that concept, the Ottawa Charter defined health as a resource for doing things—a capacity, not a state of well-being. According to this definition, health must be clearly differentiated from health status, because health has a dynamic potential for increasing or at least maintaining whatever health status (place on the spectrum) a person has. Health in this sense is a means of moving toward the positive end of the health status spectrum.

Probably more than achieving some degree of health status, people want health as a resource for doing the things they want to do. That view of health characterizes the new era of health. The goal is longevity with good function, and the challenge to health professionals is not only preventing disease and overcoming it when it occurs but also helping people to achieve that goal.

### IMPLICATIONS FOR HEALTH MEASUREMENT

What are the implications of this new third era of health and the consequent new definition of health for health measurement? Instead of just describing and analyzing the pattern of mortality, diseases, and disability, health statisticians will have to specify not only the quantitative aspects of health status but also the equal, and perhaps more important, quantitative aspects of health as the capacity for maintaining and improving health because it is a resource for living. Various elements of a person's physical, mental, and social existence make up that resource. Epidemiologists and clinicians will have to consider predictors of both functioning and disease.

As we move into the third era of health, it is useful to briefly look at both ancient and recent beliefs about health. Hippocratic thought in ancient Greece considered health to be an internal equilibrium of the 4 bodily humors: blood, phlegm, black bile, and yellow bile.<sup>14</sup> *Dyskasia*—the disturbance of that internal equilibrium—yielded disease. Because the balance between man and his environment determined

the balance of that equilibrium, factors in the environment and ways of responding to those factors profoundly affected health. Similar ideas about health emerged in ancient China.<sup>15</sup>

In 1941, Sigerist said, “Health is not simply the absence of disease: it is something positive, a joyful attitude toward life, and a cheerful acceptance of the responsibilities that life puts upon the individual.”<sup>16(p100)</sup> The WHO definition of health further inspired efforts to achieve new kinds of health measurement. Thus, Fanshel proposed 11 categories of health, 1 that encompassed the WHO notion of well-being and 10 others that ranged from dissatisfaction and discomfort through disability and coma to death.<sup>17</sup> In the Human Population Laboratory, actual measurement of health in the WHO sense of physical, mental, and social well-being was conducted with a general population survey.<sup>18–20</sup>

These past endeavors, however, focused on health as a state of positive well-being (physical, mental, and social aspects) or negative well-being (discomfort, disability, coma, etc.). The new concept advanced in the Ottawa Charter—that health is not a state of well-being but a resource for living—can be measured in its physical (e.g., body mass index [BMI]), mental (e.g., cognition), and social health dimensions (e.g., network of friends and relatives). It also can be measured in terms of health-related practices (e.g., exercise), because there is evidence that, as a category of personal characteristics, health-related practices are important resources for living that carry great influence for future health.<sup>21</sup> A schema for systematic health measurement in the third

era of health will require some consensus on not only the items to be measured but also the quantitative aspect of each item. For many items, that is well within reach.

Several items in this new kind of health measurement, especially in the physical dimension, are already being widely followed in patient care and health surveys (e.g., blood pressure, BMI, lipid level, and blood sugar). However, we usually call them risk factors and emphasize their levels of becoming abnormal, often to diagnose disease. Thus, blood pressure higher than 140 over 90 is hypertension, BMI higher than 30 is obesity, cholesterol above 200 is hypercholesterolemia, and fasting blood sugar above 126 is diabetes. We must now begin to focus on their optimal ranges, because the goal is not merely to minimize the risks for disease but to seek the maximum potential for living. Now the systematic assembly of all such details is needed for a comprehensive view of health as a resource of living. This should probably be a set of indicators rather than a single index of health.

The interpretation of the data will require establishing consensus on the quantitative range for each item that is regarded to be a part of the resource for health. Such standardization is already accepted for several items as a specification for diagnosis (e.g., blood sugar level for diabetes and hemoglobin level for anemia), but so far these are intended as indicators of a disease state. Clinicians who pursue health for their patients as I advocate here will require specific ranges that indicate an adequate reserve for everyday life. Thus, a fasting blood sugar level of 124

is too close to the diabetic state now specified as greater than 126; a healthful blood sugar range, for example, should possibly be 80 to 100. In the case of bone density, radiographers now designate a certain lack of density as osteoporosis; what I seek, however, is bone strength, not merely the absence of fragility. Therefore, we should ask the radiographers to define and measure the degree of bone density that constitutes the health of that tissue. Some might say that we must take into account age when assessing degrees of health; however, people in their later decades of life do lose some resources for living and thus, have lower health than they possessed in earlier years.

As standard health ranges are set and the relevant surveillance of the population is maintained, public health assessment and policy functions will expand to include setting population objectives for these new health indicators, just as we now set objectives for infant mortality and HIV incidence. The public health assurance function will grow to embrace community-directed activities designed to achieve the objectives for the population as a whole, and physicians will seek healthful ranges of the health indicators for their individual patients. For both public health and medical service purposes, it will be necessary to educate people as a whole about these matters; that is, we need such indicators as guidance for pursuing health—as a capacity of living—not just for avoiding disease states.

Many clinicians are moving toward the kind of health measurement I am suggesting by increasing the range of items incorporated into *comprehensive health examinations* rather than

performing the old-fashioned blood pressure cuff, stethoscopic, and manual abdominal *health check-up*. Pediatricians and obstetricians especially have moved out of the limited complaint–response practice framework toward a health maintenance system.

For public health purposes, population health surveys can begin to put together the several items of measurement necessary for each person in a survey and ultimately aggregate the health of the individuals composing the population—in an index—rather than simply determining the proportion of the population who have various blood pressure levels, cholesterol levels, BMIs, and the like. What we need is a comprehensive assembly of the items to provide a view of health as a whole resource for living—for individuals and then for the population—to guide action for health in the future. ■

#### About the Author

Lester Breslow is with the School of Public Health, University of California, Los Angeles.

Requests for reprints should be sent to Lester Breslow, MD, MPH, UCLA, School of Public Health, 650 Charles E Young Dr S, Los Angeles, CA 90095 (e-mail: breslow@ph.ucla.edu).

This article was accepted March 24, 2005.

#### Acknowledgments

I wish to acknowledge suggestions from my colleague, Robert Kaplan, and several reviewers.

#### Human Participant Protection

No protocol approval was needed for this study.

#### References

1. Terris M. The complex tasks of the second epidemiologic revolution. *J Public Health Pol.* 1983;4:8–24.
2. Keys A, ed. Coronary heart disease in seven countries. *Circulation.* 1970; 41(suppl 4):1–211.

3. Report of the Advisory Committee to the Surgeon General of the Public Health Service. *Smoking and Health.* Washington, DC: US Public Health Service; 1964. Publication No. 1103.

4. National Office of Vital Statistics, *US Public Health Services. Vital Statistics of the United States.* Vol. 1. Hyattsville, Md: 1950;209.

5. National Center for Health Statistics. *Health United States.* Hyattsville, Md: 2003.

6. Breslow L. Perspectives: the third revolution in health. *Ann Rev Public Health.* 2004;25:xiii–xviii.

7. World Health Organization (WHO), European Regional Office. *Ottawa Charter for Health Promotion.* Copenhagen, Denmark: WHO; 1986.

8. WHO. *Basic Documents.* 35th ed. Geneva, Switzerland: WHO; 1985.

9. WHO. *Measurement of Levels of Health.* Geneva, Switzerland: WHO; 1957.

10. Moriyama IM. Problems in the measurement of health status. In: Sheldon E, Moore W, eds. *Indicators of Social Change.* New York, NY: Russell Sage Foundation; 1968.

11. Chen, MK, Bryant, BE. The measurement of health—a critical and selective overview. *Int J Epidemiology.* 1975; 4:257–264.

12. Kaplan RM, Bush JW, Berry CC. Health status index-category rating versus magnitude estimation for measuring levels of well-being. *Med Care.* 1979;17: 501–525.

13. Breslow L. A quantitative approach to the World Health Organization definition of health: physical, mental, and social well-being. *Int J Epidemiol.* 1972; 4:347–355.

14. Sigerist HE. *Civilization and Disease.* Ithaca, NY: Cornell University Press; 1943.

15. Bowers Z. *Medicine and Society in China.* Philadelphia, Pa: Josiah May Jr Foundation; 1974.

16. Sigerist HE. *Medicine and Human Welfare.* New Haven, Conn: Yale University Press; 1941.

17. Fanshel S. A meaningful measure of health for epidemiology. *Int J Epidemiol.* 1972;1:319–337.

18. Belloc N, Breslow L, Hochstim J. Measurement of physical health in a general population survey. *Am J Epidemiol.* 1971;93:328–336.

19. Berkman PL. Measurement of mental health in a general population survey. *Am J Epidemiol.* 1971;94: 104–111.

20. Renne K. Measurement of social health in a general population survey. *Soc Sci Res.* 1974;1:25–44.

21. Breslow L, Breslow N. Health practices and disability: some evidence from Alameda County. *Prev Med.* 1993;22: 86–95.

**This article has been cited by:**

1. Admassu N. Lamu, Jan Abel Olsen. 2018. Yes, health is important, but as much for its importance via social life: The direct and indirect effects of health on subjective well-being in chronically ill individuals. *Health Economics* 27:1, 209-222. [[Crossref](#)]
2. Aditya Jain, Stavroula Leka, Gerard I. J. M. Zwetsloot. Work, Health, Safety and Well-Being: Current State of the Art 1-31. [[Crossref](#)]
3. Jerome Bickenbach. WHO's Definition of Health: Philosophical Analysis 961-974. [[Crossref](#)]
4. Pascale Mantoura, Marie-Claude Roberge, Louise Fournier. 2017. Un cadre de référence pour soutenir l'action en santé mentale des populations. *Santé mentale au Québec* 42:1, 105. [[Crossref](#)]
5. Y Piri, A Zeinali. 2016. Relationship between Perceived Social Support, Social Capital and Quality of Life with Job Burnout among Nurses. *Iran Journal of Nursing* 29:103, 13-23. [[Crossref](#)]
6. Pascale Lehoux, Federico Roncarolo, Robson Rocha Oliveira, Hudson Pacifico Silva. 2016. Medical innovation and the sustainability of health systems: A historical perspective on technological change in health. *Health Services Management Research* 29:4, 115-123. [[Crossref](#)]
7. J. K. Preskitt, K. S. Menear, S. S. Goldfarb, N. Menachemi. 2015. Wellness among US adolescents ages 12-17 years. *Child: Care, Health and Development* 41:6, 1207-1215. [[Crossref](#)]
8. Craig M. Becker, Mary A. Glascoff, William Michael Felts, Christopher Kent. 2015. Adapting and Using Quality Management Methods to Improve Health Promotion. *EXPLORE: The Journal of Science and Healing* 11:3, 222-228. [[Crossref](#)]
9. K.S. Menear, J.K. Preskitt, S.S. Goldfarb, N. Menachemi. 2015. Correlates of wellness among youth with functional disabilities. *Disability and Health Journal* 8:2, 223-230. [[Crossref](#)]
10. Peter D Wong, Malini G Dave, Trisha Tulloch, Mark L Feldman, Elizabeth L Ford-Jones, Patricia C Parkin, Adrianna Tetley, Rosemary G Moodie. 2015. Community health centres: Potential opportunities for community paediatrics. From interprofessional clinical care to board governance. *Paediatrics & Child Health* 20:1, 12-14. [[Crossref](#)]
11. Peipei Song, Yu Chen. 2015. Public policy response, aging in place, and big data platforms: Creating an effective collaborative system to cope with aging of the population. *BioScience Trends* 9:1, 1-6. [[Crossref](#)]
12. Neal Halfon, Peter Long, Debbie I. Chang, James Hester, Moira Inkelas, Anthony Rodgers. 2014. Applying A 3.0 Transformation Framework To Guide Large-Scale Health System Reform. *Health Affairs* 33:11, 2003-2011. [[Crossref](#)]
13. Neal Halfon, Kandyce Larson, Michael Lu, Ericka Tullis, Shirley Russ. 2014. Lifecourse Health Development: Past, Present and Future. *Maternal and Child Health Journal* 18:2, 344-365. [[Crossref](#)]
14. Theodore H. Tulchinsky, Elena A. Varavikova. Measuring, Monitoring, and Evaluating the Health of a Population 91-147. [[Crossref](#)]
15. LM Anderson, JB Moore, BM Hayden, CM Becker. 2014. Test-retest reliability of the Salutogenic Wellness Promotion Scale (SWPS). *Health Education Journal* 73:1, 101-108. [[Crossref](#)]
16. Rodolfo J. Stusser, Richard A. Dickey. 2013. Quality and Cost Improvement of Healthcare via Complementary Measurement and Diagnosis of Patient General Health Outcome Using Electronic Health Record Data: Research Rationale and Design. *Journal of Medical Systems* 37:6. . [[Crossref](#)]
17. Rosemarie Kobau, Carla Bann, Megan Lewis, Matthew M Zack, Angela M Boardman, Renee Boyd, Kim C Lim, Tommy Holder, Anastacia KL Hoff, Cecily Luncheon, William Thompson, Willi Horner-Johnson, Richard E Lucas. 2013. Mental, social, and physical well-being in New Hampshire, Oregon, and Washington, 2010 Behavioral Risk Factor Surveillance System: implications for public health research and practice related to Healthy People 2020 foundation health measures on well-being. *Population Health Metrics* 11:1. . [[Crossref](#)]
18. Katarzyna Byrka, Florian G. Kaiser. 2013. Health performance of individuals within the Campbell paradigm. *International Journal of Psychology* 48:5, 986-999. [[Crossref](#)]
19. Blaine Reeder, Ellen Meyer, Amanda Lazar, Shomir Chaudhuri, Hilaire J. Thompson, George Demiris. 2013. Framing the evidence for health smart homes and home-based consumer health technologies as a public health intervention for independent aging: A systematic review. *International Journal of Medical Informatics* 82:7, 565-579. [[Crossref](#)]
20. Craig M. Becker, Patricia Rhynders. 2013. It's time to make the profession of health about health. *Scandinavian Journal of Public Health* 41:1, 1-3. [[Crossref](#)]
21. Diane E. Mack, Philip M. Wilson, Katie E. Gunnell, Jenna D. Gilchrist, Kent C. Kowalski, Peter R.E. Crocker. 2012. Health-Enhancing Physical Activity: Associations with Markers of Well-Being. *Applied Psychology: Health and Well-Being* 4:2, 127-150. [[Crossref](#)]

22. Nady el-Guebaly. 2012. The Meanings of Recovery From Addiction. *Journal of Addiction Medicine* 6:1, 1-9. [[Crossref](#)]
23. José L. Pais-Ribeiro. 2012. Exploration of an Agentic Construct that Impacts Health Behaviors in General Population. *Psychology* 03:09, 698-701. [[Crossref](#)]
24. Blaine Reeder, George Demiris. 2011. Smart homes and novel indicators to inform an evidence-based population health intervention for aging in place and design of a community health registry. *Emerging Health Threats Journal* 4:0. . [[Crossref](#)]
25. Andrew Pleasant, Julie McKinney. 2011. Coming to consensus on health literacy measurement: An online discussion and consensus-gauging process. *Nursing Outlook* 59:2, 95-106.e1. [[Crossref](#)]
26. Mirjana Arandjelovic. 2011. A need for holistic approach to the occupational health developing (in Serbia). *International Journal of Occupational Medicine and Environmental Health* 24:3. . [[Crossref](#)]
27. Rosemarie Kobau, Joseph Sniezek, Matthew M. Zack, Richard E. Lucas, Adam Burns. 2010. Well-Being Assessment: An Evaluation of Well-Being Scales for Public Health and Population Estimates of Well-Being among US Adults. *Applied Psychology: Health and Well-Being* 2:3, 272-297. [[Crossref](#)]
28. Jürgen Rehm, Ulrich Frick. 2010. Valuation of health states in the US study to establish disability weights: lessons from the literature. *International Journal of Methods in Psychiatric Research* 19:1, 18-33. [[Crossref](#)]
29. Adam Buciński, Jerzy Romaszko, Ryszard Targoński, Leszek Cichowski, Ewa Romaszko, Andrzej Zakrzewski, Małgorzata Wnuk. 2010. Evaluation of selected lipid parameters and blood pressure in ethnically-homogenous population of middle-aged persons, depending on gender, age and body mass. *Open Medicine* 5:4. . [[Crossref](#)]
30. Deanna L. Williamson, Jeff Carr. 2009. Health as a resource for everyday life: advancing the conceptualization. *Critical Public Health* 19:1, 107-122. [[Crossref](#)]
31. Kakoli Roy, Anne C. Haddix, Robin M. Ikeda, Cecilia W. Curry, Benedict I. Truman, Stephen B. Thacker. 2009. Monitoring Progress toward CDC's Health Protection Goals: Health Outcome Measures by Life Stage. *Public Health Reports* 124:2, 304-316. [[Crossref](#)]
32. Alexandre B. Laudet, Jeffrey B. Becker, William L. White. 2009. Don't Wanna Go Through That Madness No More: Quality of Life Satisfaction as Predictor of Sustained Remission from Illicit Drug Misuse. *Substance Use & Misuse* 44:2, 227-252. [[Crossref](#)]
33. Craig Becker, Lauren Whetstone, Mary Glascoff, Justin B. Moore. 2008. Evaluation of the Reliability and Validity of an Adult Version of the Salutogenic Wellness Promotion Scale (SWPS). *American Journal of Health Education* 39:6, 322-328. [[Crossref](#)]
34. Elizabeth Lee Ford-Jones, Robin Williams, Jane Bertrand. 2008. Social paediatrics and early child development: Part 1. *Paediatrics & Child Health* 13:9, 755-758. [[Crossref](#)]
35. Alexandre Laudet. 2008. Editorial. *Substance Use & Misuse* 43:12-13, 1681-1684. [[Crossref](#)]
36. Craig Becker, Christyn L. Dolbier, Thomas W. Durham, Mary A. Glascoff, Troy B. Adams. 2008. Development and Preliminary Evaluation of a Positive Health Scale. *American Journal of Health Education* 39:1, 34-41. [[Crossref](#)]
37. 2007. What is recovery? A working definition from the Betty Ford Institute. *Journal of Substance Abuse Treatment* 33:3, 221-228. [[Crossref](#)]
38. James E. Rohrer, Ahmed Arif, Anne Denison, Rodney Young, Steve Adamson. 2007. Overall self-rated health as an outcome indicator in primary care. *Journal of Evaluation in Clinical Practice*, ahead of print070821070410006-???. [[Crossref](#)]
39. James E. Rohrer, Stephen P. Merry, Francisco Lopez-Jimenez, Steven C. Adamson, Laurie Wilshusen. 2007. A Patient-Centered Decision Rule for Referral of Patients to Weight-Loss Programs. *Quality Management in Health Care* 16:3, 250-255. [[Crossref](#)]
40. Donna Allen, Deb Carlson, 'Chelle Ham. 2007. Well-Being: New Paradigms of Wellness—Inspiring Positive Health Outcomes and Renewing Hope. *American Journal of Health Promotion* 21:3, 1-12. [[Crossref](#)]
41. Barbara K Redman. 2006. Ethically problematic assumptions regarding patient self management and barriers to improved outcomes. *Expert Review of Pharmacoeconomics & Outcomes Research* 6:5, 489-494. [[Crossref](#)]
42. Jacqueline L. Angel, Ronald J. Angel. 2006. Minority Group Status and Healthful Aging: Social Structure Still Matters. *American Journal of Public Health* 96:7, 1152-1159. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
43. Amirhosein Ghaffarianhoseini, Ali Ghaffarianhoseini, John Tookey, Hossein Omrany, Anthony Fleury, Nicola Naismith, Mahdiar Ghaffarianhoseini. The Essence of Smart Homes 334-376. [[Crossref](#)]